

---

# Eye Avoidance in Young Children With Autism Spectrum Disorder Is Modulated by Emotional Facial Expressions

Qiandong Wang and Li Lu  
Peking University

Qiang Zhang  
Sun Yat-sen University

Fang Fang  
Peking University

Xiaobing Zou  
Sun Yat-sen University

Li Yi  
Peking University

Individuals with autism spectrum disorder (ASD) exhibit a reduced duration of eye contact compared with typically developing (TD) individuals. This reduced eye contact has been theorized to be a strategy to relieve discomfort elicited by direct eye contact (Tanaka & Sung, 2016). Looking at threatening facial expressions may elicit more discomfort and consequently more eye avoidance in ASD individuals than looking at nonthreatening expressions. We explored whether eye avoidance in children with ASD is modulated by the social threat level of emotional expressions. In this study, 2- to 5-year-old children with and without ASD viewed faces with happy, angry, sad, and neutral expressions, while their eye movements were recorded. We observed the following: (a) when confronted with angry faces, the children with ASD fixated less on the eyes than did TD children, persistently across time; (b) the group differences in the overall eye-looking time were rarely found for happy, neutral, and sad faces; (c) the ASD group showed eye avoidance for neutral faces between 1,000 ms and 2,900 ms after the stimulus onset. Additionally, both groups spent more time looking at the angry faces than the faces showing other emotions. Considering that the children with ASD spent less time looking at the eyes of the angry faces than other emotional faces, the results suggest a combination of vigilance to threatening faces and an avoidance of the eyes in children with ASD. Our study not only extends the gaze aversion hypothesis but also has implications for the treatment and screening of ASD.

## General Scientific Summary

Previous studies have indicated that individuals with ASD exhibit an eye-avoidance pattern when scanning faces. This study observed that this pattern is modulated by facial expressions; the eye-avoidance pattern in young children with ASD is specific to threatening facial expressions, which suggests that eye avoidance may help children with ASD alleviate discomfort elicited by threatening facial expressions.

**Keywords:** autism spectrum disorder, face scanning, eye avoidance, eye movement, emotional expressions

**Supplemental materials:** <http://dx.doi.org/10.1037/abn0000372.supp>

---

Qiandong Wang, Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University; Li Lu, School of Psychological and Cognitive Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University; Qiang Zhang, Southern China Research Center of Statistical Science, School of Mathematics, Sun Yat-sen University; Fang Fang, Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, School of Psychological and Cognitive Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University; Xiaobing Zou, The Third Affiliated Hospital of Sun Yat-sen University, Sun Yat-sen University; Li Yi, School of Psychological and Cognitive

Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University.

This study was supported by National Natural Science Foundation of China (31571135, 11771462) and Beijing Municipal Science and Technology Commission (Z171100000117015). The authors are grateful to Lisa Joseph, Nikki Greenwood, Bella Chan, Jiale Li, Jiao Li, Jianling Tang, Dejun Shi, Sinong Chen, and children and parents who participated in our study.

Correspondence concerning this article should be addressed to Li Yi, School of Psychological and Cognitive Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China 100871. E-mail: [yilipku@pku.edu.cn](mailto:yilipku@pku.edu.cn)

Autism spectrum disorder (ASD) is a pervasive neurodevelopmental disorder characterized by impairments in social interaction and communication, as well as by the presence of repetitive or stereotypical behaviors (American Psychiatric Association, 2013). Along with these social deficits, individuals with ASD have been found to have impairments in facial identity discrimination (Weigelt, Koldewyn, & Kanwisher, 2012) and emotion recognition (Rump, Giovannelli, Minshew, & Strauss, 2009; Uljarevic & Hamilton, 2013). People with ASD also exhibit atypical neural responses to faces (Dalton et al., 2005; Pierce, Müller, Ambrose, Allen, & Courchesne, 2001) and atypical face-scanning patterns (for a review, see Falck-Ytter & von Hofsten, 2011). Particularly, eye-tracking studies have repeatedly found that people with ASD spend less time looking at others' eyes than typically developing (TD) counterparts (e.g., Jones & Klin, 2013; Pelphrey et al., 2002; Yi et al., 2013). This is referred to as the "eye avoidance" looking pattern in ASD (Tanaka & Sung, 2016). These findings are consistent with clinical observations that individuals with ASD have limited eye contact with others (Adrien et al., 1993; however, see later).

The mechanisms underlying this atypical eye-gaze in ASD have attracted increased research interest in the past decade. Some researchers have proposed that individuals with ASD perceive direct eye contact as socially threatening, and thus actively avoid looking at others' eyes to relieve the uncomfortable feelings elic-

are eye,ng loon

ingffntentnlgactedthreng socingffntentng  
to styhe The

with time. The former would indicate a strong, consistent form of eye avoidance, suggesting that children with ASD cannot habituate to social threats, and thus continue to avoid looking at the eyes. The latter, on the other hand, would suggest a mild form of eye avoidance, perhaps representing their habituation to the threat over time. We expected to see different eye-avoidance patterns when children with ASD were processing faces with different facial expressions—more specifically, we expected stronger forms of eye avoidance in more threatening facial expressions (e.g., anger). The temporal-course analysis was intended to reveal the nuances in children's responses to different facial expressions over time, which is of importance to our understanding of emotional face processing in ASD. Such nuanced responses to different facial expressions could also distinguish ASD from other disorders (e.g., social anxiety), and might have implications for the development of training programs targeted at improving eye contact in individuals with ASD.

## Method

### Participants and Ethical Considerations

We recruited thirty 2- to 5-year-old Chinese children with ASD, who were diagnosed by pediatric psychiatrists according to the diagnostic criteria for ASD in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-V; American Psychiatric Association, 2013)*. Children with ASD were further assessed by using the Autism Diagnostic Observation Schedule (ADOS; Lord et al., 2000). Three children were excluded from the study as they did not meet diagnostic criteria based on the ADOS. Ultimately, 27 children with ASD (four girls,  $M_{\text{age}} = 40.3$  months,  $SD_{\text{age}} = 10.3$  months) participated in our study (see Table 1). We also recruited 31 age-matched TD children (five girls,  $M_{\text{age}} = 41.2$  months,  $SD_{\text{age}} = 9.2$  months). The TD children were recruited at their routine wellness examinations at local hospitals, and they did not exhibit any signs of ASD or other developmental issues. The present protocol (protocol number: 2016-03-03e) was approved by the Committee for Protecting Human and Animal Subjects at School of Psychological and Cognitive Sciences at Peking University, China. We obtained all children's verbal assent and their parents' written consent before commencing with the experiment.

## Materials

The faces used in the present study included four neutral, four happy, four angry, and four sad female faces. We drew these face photos from the Chinese Facial Affective Picture System (CFAPS; Gong, Huang, Wang, & Luo, 2011; Wang & Luo, 2005) stimulus set. All facial images (width, 260 pixels; height, 300 pixels) were frontal views and were rendered in gray scale, with hair eliminated (see Figure 1). We used female faces as past studies indicated children are more familiar with female adult faces than male faces (Quinn, Yahr, Kuhn, Slater, & Pascalis, 2002).

## Procedure

Children sat approximately 50 cm away from the screen. Parents were asked to sit behind them (out of view of the eye tracker) and to remain quiet. We used a Tobii T60 eye tracker (TobiiTech, Stockholm, Sweden), an integrated eye tracker and presentation desktop system with 60 Hz sample rate and  $1,024 \times 768$  pixel resolution, to record the gaze data. The Tobii Studio 1.5 software (TobiiTech, Stockholm, Sweden) was used to control the stimulus presentation. We used the Tobii's five-point calibration method for every child. We accepted the calibration only if all five points for the two eyes were caught by the Tobii, with only small error vectors (smaller than 0.5 degree of visual angle).

A trial began with a black screen with a white cross at the center of the monitor (approximately at the nose region relative to the face) for 500 ms. One face was later displayed for 5,000 ms at the center of the computer screen, and the children were instructed to look at it freely. The 16 pictures, four in each facial expression, were randomly presented (i.e., a new random order for each participant). The interstimulus interval between the face presentations was 500 ms with a white cross presented at the center of the monitor. After every two trials, a cartoon video was presented for 10 s to maintain the child's attention.

## Data Analysis

We defined four areas of interest (AOIs) for each face: the right eye, the left eye, the mouth, and the whole face (see Figure 1). We defined the eye region as described in earlier studies (eyebrows not

Table 1  
Mean (SD) Scores on the Autism Diagnostic Observation Schedule in Children With ASD

Module (Number of participants)	Domain	M (SD)
Module 1 (N = 16)	Communication	6.3 (2.4)
	Social interaction	8.9 (3.0)
	Communication & social interaction	15.2 (4.7)
	Play	2.1 (1.2)
	Stereotyped behaviors	2.3 (1.1)
Module 2 (N = 11)	Communication	5.8 (1.7)
	Social interaction	8.3 (2.1)
	Communication and social interaction	14.1 (3.5)
	Imagination/creativity	1.0 (.5)
	Stereotyped behaviors	1.5 (1.0)

Note. Sixteen out of 27 children with ASD with low-level language (no speech to simple phrases) were conducted with Module 1, and 11 out of 27 children with ASD with higher-level of language (use phrase speech but not yet achieve verbal fluency) were conducted with Module 2.

included; e.g., Yi et al., 2013), and we combined the time spent at the left and right eyes and used that result as the time spent at the eyes, or eye-*looking* time. The “whole face” region included the area within the face contour. Because the current study primarily focused on the eyes, supplementary analysis of the nose and the nonfeatural face region (face areas excluding the eyes, nose, and mouth) can be found in the [online supplementary material \(Figure S1 and Table S1\)](#). We also computed the Eye-Mouth Index (the difference between eye-*looking* time and mouth-*looking* time), and have reported it in the [online supplementary material \(Table S2\)](#).

We used the gaze data (i.e., sample data) rather than the fixation data as an analytical unit. We first computed the total face-*looking* time by summing all gaze durations on the whole face for each expression. Next, we calculated the proportional *looking* time on the AOIs of the eyes and mouth, respectively, by dividing the total *looking* time on each AOI by the total *looking* time on the whole face for each expression. We excluded the trials in which participants spent no time *looking* at the face, to avoid invalid denominators; the average number of invalid trials is listed in [Table 2](#). At least one valid trial for each expression was necessary to be admitted into the study. According to these exclusion criteria, one child with ASD did not have any valid trials when scanning the sad faces and thus was excluded from any analysis including sad faces. To examine whether there was a difference in the number of invalid trials across groups and expressions, we conducted a 2 (Group)  $\times$  4 (Expression) repeated measures ANOVA on the invalid trial number, and found a significant main effect of group, suggesting more invalid trials in the ASD than the TD groups, and an effect of expression (see [Table 2](#)).

To test whether eye avoidance in ASD was modulated by different expressions, we used ANOVA and *t* tests (two-tailed) to test our hypothesis, and used false discovery rate (FDR) adjustment for multiple comparisons to control for Type I error.

We further adopted two temporal-course analyses to examine how the eye-avoidance pattern in ASD changed over time, and how the eye-*looking* time changed over time for each group. First, we examined the temporal course of the scanning 0(tempora.8(exprmJ1cr8(exp

for each cluster until none of the maximum values in any cluster is significant.

The two temporal-course analyses are described in more detail in the [online supplementary material](#).

## Results

### Total Looking Time on the Whole Face

We first examined group differences regarding total looking time on the whole face using a 2 Group (ASD and TD groups)  $\times$  4 Expression (Angry, Neutral, Happy, and Sad) repeated-measures ANOVA. As shown in [Figure 2](#) and [Table S3](#), only the main effect of Expression was significant,  $F(3, 165) = 31.99, p < .001, \eta_p^2 = .37, 90\% \text{ CI } [.26, .44]$ . Post hoc pairwise  $t$  tests (after FDR correction) revealed that children looked longer at angry than neutral, happy, and sad faces,  $t(57) = 4.43, p < .001$ , Cohen's  $d = 0.58, 95\% \text{ CI } [0.30, 0.86]$ ;  $t(57) = 7.05, p < .001$ , Cohen's  $d = 0.93, 95\% \text{ CI } [0.61, 1.23]$ ;  $t(56) = 7.48, p < .001$ , Cohen's  $d = 0.99, 95\% \text{ CI } [0.67, 1.31]$ , respectively. Children also looked longer at neutral faces than happy and sad faces,  $t(57) = 3.51, p = .001$ , Cohen's  $d = 0.46, 95\% \text{ CI } [0.19, 0.73]$ ;  $t(56) = 4.53, p < .001$ , Cohen's  $d = 0.60, 95\% \text{ CI } [0.32, 0.88]$ , respectively. The difference in face-looking time between happy and sad faces was not significant,  $t(56) = 1.87, p = .067$ , Cohen's  $d = 0.25, 95\% \text{ CI } [-0.02, 0.51]$ . There was no significant main effect of group,  $F(1, 55) = 0.22, p = .64, \eta_p^2 < .01, 90\% \text{ CI } [.00, .07]$ , or the Group  $\times$  Expression interaction,  $F(3, 165) = 2.31, p = .087, \eta_p^2 = .04, 90\% \text{ CI } [.00, .09]$ , indicating a similar amount of time looking at the whole face for the two groups. The same analyses were applied to the total time spent looking at the screen, and results were very similar to those of total time spent looking at the whole face (see [Table S3](#) and [Figure S2](#) for more details).

### Proportional Eye-Looking Time

We examined whether the ASD and TD groups showed different proportional looking time focused on the eyes for different

facial expressions across total viewing time. A repeated-measures ANOVA was conducted on the proportional eye-looking time, with expression as the within-subject variable, and group as the between-subjects variable. The results are shown in [Figure 3A](#) and [Table S4](#).

Both main effects of group and expression on the proportional eye-looking time were significant,  $F(1, 55) = 5.09, p = .028, \eta_p^2$




Figure 3. Proportional looking time on the eyes and mouth of different emotional faces of the ASD and the TD groups (error bars denote standard errors; \*\*\* denotes  $p < .001$ ). See the online article for the color version of this figure.

and neutral versus sad,  $t(30) = -2.20$ ,  $p = .07$ , Cohen's  $d = 0.40$ , 95% CI [0.03, 0.76].

### Temporal-Course Analysis

Aside from the analyses that collapsed all time spent looking at the eyes across the total viewing time, we further examined the temporal course of the eye-looking time to determine when the eye avoidance appeared and how it changed over time. The results are presented in Figure 4. For the angry faces, the ASD group spent less time looking at the eyes than the TD group in almost all epochs. For the neutral faces, significant group differences in eye-looking time existed between about 1,000 ms and 2,900 ms after the face appeared, and the two groups looked at the eyes similarly in other epochs. For happy faces, the two groups displayed similar eye-looking time in almost all epochs with the exception of a short interval, 616 ms–682 ms, after the face appeared. For sad faces, the two groups spent similar time on the eyes in all epochs.

Additionally, we examined how eye-looking time changed over time for each group and expression respectively. Change-points found by the BDCP analysis are illustrated in Figure 5. For angry faces, only one change-point was found for the ASD group, and no change-points were found for the TD group. For other expressions, multiple change-points were found, suggesting that proportional eye-looking time for both groups oscillated across time—they increased their attention to the eyes gradually and then gradually decreased their attention to the eyes, and repeated this procedure across time. The oscillation amplitudes, however, changed across time.

Combining these two temporal-course approaches, we reached the following conclusions. (a) For angry faces, the ASD group displayed consistently shorter proportional eye-looking time across time relative to the TD group. (b) For neutral faces, the eye-avoidance pattern of ASD occurred around 1,000 ms after the onset of the face, and lasted for around 2,000 ms (see Figure 4). Due to both the increased proportional eye-looking time of the

ASD group and the decreased proportional eye-looking time of the TD group between 1,500 ms and 3,000 ms (see Figure 5), the group difference disappeared after 2,900 ms. (c) For the happy and the sad faces, the proportional eye-looking time was very similar in the two groups across time.

### Proportional Mouth-Looking Time

A 4 (Expression)  $\times$  2 (Group) repeated-measures ANOVA on the proportional mouth-looking time found no significant main effect of group,  $F(1, 55) = 0.05$ ,  $p = .83$ ,  $\eta_p^2 < .01$ , 90% CI [.00, .04]; expression,  $F(3, 165) = 2.03$ ,  $p = .11$ ,  $\eta_p^2 = .04$ , 90% CI [.00, .08]; or their interaction,  $F(3, 165) = 1.35$ ,  $p = .26$ ,  $\eta_p^2 = .02$ , 90% CI [.00, .06]. That is, the two groups looked at the mouth similarly for all four expressions (Figure 3B and Table S4).

### Discussion

In the present study, we employed eye-tracking to examine eye avoidance in young children with ASD when processing facial expressions. We observed that the hypothesized eye avoidance in ASD was most prominent for threatening facial expressions (i.e., angry faces), which is consistent with the gaze aversion hypothesis that considers eye avoidance as a strategy to relieve discomfort elicited by social threat (Hutt & Ounsted, 1966; Kliemann et al., 2010; Tanaka & Sung, 2016). The results of the within-group comparisons further revealed that the children with ASD looked at the eyes of the angry faces less than the neutral, happy, and sad faces, while TD children looked at the eyes of angry and sad faces more than happy faces. Furthermore, our temporal-course analysis revealed for angry faces, children with ASD showed the eye-avoidance pattern relative to the TD children in almost all time epochs; for neutral faces, the eye avoidance in children with ASD began about 1,000 ms after the onset of the face, and lasted to 2,900 ms. For the happy and sad faces, group differences were rarely found at any time. These findings could not be explained by the group differences in the overall attention distribution, given the similar face-looking time between the two groups.






Figure 4. Proportional eye-looking time of the ASD and the TD groups for different expressions over time (shaded area indicates standard errors). Gray shade illustrates the cluster of time epochs when the group differences of eye-looking time are significant. See the online article for the color version of this figure.

Our comparative findings, in which the ASD group showed reduced proportional eye-looking time of angry faces, but not of sad faces, suggest that the eye avoidance of ASD has a greater likelihood of being associated with more socially threatening stimuli, rather than with negativity as a whole. Further evidence came from the analyses that compared the face-scanning patterns between different expressions for each group. The TD children looked at the eyes of the angry and sad faces more than those of the happy faces. Considering that expressive information of negative expressions is, for the most part, conveyed by the upper half of the face (Eisenbarth & Alpers, 2011; Schurkin et al., 2014; Smith, Cottrell, Gosselin, & Schyns, 2005), our finding suggests that TD children are very sensitive to this nuance among expressions. However, such sensitivity was absent in the children with ASD, who looked at the eyes of the angry faces less than the neutral, happy, and sad faces, further suggesting that eye avoidance in ASD is specific to socially threatening expressions. These interexpression comparisons also suggest that the decreased eye-looking time of children with ASD for angry faces is due to their atypical face processing, but not to their decreased physical salience

for the eyes of angry faces, given that the TD children spent the most time looking at the eyes of angry faces.

Despite their eye avoidance responses toward angry faces, the children with ASD, similar to TD children, looked more at the whole faces with threatening emotional expressions than those with non-threatening expressions, which is consistent with several previous studies (Hall, Hutton, & Morgan, 2012; Perez-Edgar et al., 2017). Faces displaying threatening emotional expressions, such as anger and fear, have been described to evoke vigilance responses from individuals (Green, Williams, & Davidson, 2003), leading to longer looking time for these faces. In fact, infants as young as 7 months already show difficulty in disengaging from threatening faces (Peltola, Lepänen, Vogel-Farley, Hietanen, & Nelson, 2009). It should be noted that although the children with ASD paid more attention to the angry faces than to the nonthreatening faces, they looked less at the inner features of the faces (especially the eyes). That is, the children with ASD did not concentrate on the eyes, nose, and mouth of the angry faces, but distributed their gaze to a greater extent (more than half of the time), over the nonfeatural face region (Table S1), strongly indi-




Figure 5. Change-point analysis. Change-points are marked by square for the ASD group and diamond for the TD group. Average proportional eye-looking time before, after, or between change points are marked by dash line for the ASD group and dot line for the TD group. The solid lines represent proportional eye-looking time across time. See the online article for the color version of this figure.

cating a complete avoidance of the core facial features specific to angry faces. This looking style might be a combination of vigilance to threatening faces and avoidance to the core features of such faces, especially the eyes, which convey the most social and threatening information (Eisenbarth & Alpers, 2011; Farabee, Holcom, Ramsey, & Cole, 1993; Smith et al., 2005; Wieser, Pauli, Alpers, et al., 2009). While further research is certainly needed, such a special attentional style is obviously meaningful to people with ASD. For example, being more vigilant to threatening faces can help people with ASD detect important social cues; also, avoiding eye contact can alleviate discomfort elicited by threatening facial expressions. Notably, this looking pattern reflects a *spatial* distribution of visual attention, and is different from the vigilance-avoidance looking pattern found in people with social anxiety (Holas et al., 2014; Wieser, Pauli, Weyers, et al., 2009), which, revealed by the temporal-course analyses, represents a *temporal* change of looking pattern toward social threat (initially enhanced and then subsequently reduced looking time).

With regard to the temporal-course analysis, as expected, the children with ASD demonstrated a strong form of eye avoidance by

showing eye avoidance for the threatening facial expression (anger) persistently over the presentation time. For the neutral faces, although the group difference of the overall eye-looking time was not significant, the temporal-course analysis revealed that eye avoidance did appear between 1,000 ms and 2,900 ms after the face onset. After that, the group difference disappeared, due to both increased eye-looking time of the ASD group and decreased eye-looking time of the TD group. This eye-avoidance pattern confirms the mild form of eye avoidance in line with our hypothesis. Neutral expressions are often found to be confused with negative and threatening faces by both TD participants and participants with ASD (Eack, Mazefsky, & Minshew, 2015; Juth, Lundqvist, Karlsson, & Ohman, 2005; Lee, Kang, Park, Kim, & An, 2008). Thus, the decreased eye-looking with neutral faces in children with ASD may still reflect their tendency to avoid the threat-elicited discomfort posed by the eyes of neutral facial expressions. Given the lower degree of threat of neutral faces relative to angry faces, children with ASD showed a mild form of the eye-avoidance pattern that eclipsed over time. This can be attributed to



their habituation or their reduced perception of the threat of neutral faces over time.

Our temporal-course analysis has additional implications. First, most previous research measured gazing behavior across the whole stimulus-presenting time without evaluating the exact temporal course of attention allocation to the eyes (e.g., De Wit et al., 2008; Yi et al., 2013). Our results suggest that attention allocation may depend on the length of the stimulus presentation time. Thus, it may provide insights into the inconsistent findings in previous studies regarding emotional face-scanning in ASD. Second, this temporal-course measure may help disentangle the abnormal attentional processes in different psychiatric disorders. It has been shown that people with social anxiety avoid looking at the eyes as well (Horley et al., 2003; Wang, Hu, Short, & Fu, 2012); however, they also show an initial vigilance (e.g., enhanced eye-looking time) followed by avoidance of social threat (e.g., reduced eye-looking time), which has been interpreted as evidence of a vigilant-avoidant attentional bias (Boll et al., 2016; Wieser, Pauli, Weyers, et al., 2009). In contrast, in our study children with ASD did not show enhanced eye-looking time toward the threatening expressions. Our findings suggest the feasibility of developing an algorithm to classify individuals with ASD and those with social anxiety simply based on their nuanced temporal courses of eye-looking patterns.

We failed to demonstrate the longer mouth-looking time in the ASD group relative to the TD group found by previous eye-tracking studies using dynamic videos (Jones, Carr, & Klin, 2008; Klin, Jones, Schultz, Volkmar, & Cohen, 2002). It is possible that when presented with a dynamic stimulus and in a communicative situation, individuals with ASD may be more attracted by the movements of the mouth, which is compatible with their preserved ability to use visual information from the mouth for speech-related processing (Klin et al., 2002). Previous studies using static faces have found similar results as the current study by showing comparable or even reduced mouth-looking time in ASD as compared with TD children (e.g., Fedor et al., 2017; Yi et al., 2013). The similar mouth-looking time in ASD and TD groups found in our study has ruled out the possibility that eye avoidance in ASD is driven by increased attention to, or interest in, the visual and communicative information conveyed by the mouth.

Several main considerations emerge from these findings. First, similar to most previous studies on face processing in ASD, we used static photos of emotional expressions. The generalization of our conclusions should be examined in future studies using more ecologically valid stimuli, perhaps even live stimuli (e.g., interpersonal interactions between the participants and an experimenter) using head-mounted eye-tracking equipment. Second, the lack of measurement of IQ or developmental level of the current sample is another limitation of the current study. Although a recent meta-analysis indicated no impact of IQ on gaze abnormalities in ASD (Frazier et al., 2017), from a scientific point of view it is crucial to match this variable between groups to ensure that the observed group differences were due to the diagnostic status rather than to developmental levels. To better illustrate the relationship between IQ and the eye-avoidance pattern in ASD, future work could study the gaze patterns of children with ASD with a broad range of IQ. Third, future investigations could explore the link between eye contact and arousal in young children with ASD by

recording skin conductance and eye movements simultaneously. Future studies could also use the eye-tracking technique combined with questionnaires about anxiety and neuroimaging techniques to further explore the relationships between anxiety and eye avoidance in ASD. Fourth, future studies should examine the effects of other types of threatening facial features (e.g., aggressiveness and gender, Carré, McCormick, & Mondloch, 2009; Renzi, Tagliaferri, & Boehringer, 2014) on eye-looking patterns in ASD. For example, our study only used female faces, and it is unclear whether male faces, which could be perceived as more threatening than female faces (Renzi et al., 2014), elicit a stronger eye-avoidance pattern. Additionally, we could cue children's attention to the eyes, as Moriuchi, Klin, and Jones (2017) did, but with faces of different expressions to test whether facial expressions would modulate the extent of "flight from the eyes" in ASD (Moriuchi, Klin, & Jones, 2017). Fifth, the atypical face-scanning pattern was used as a potential marker for screening infants and children with ASD in previous studies (e.g., Jones & Klin, 2013; Liu, Li, & Yi, 2016). The findings of the present study imply that angry faces may be a more powerful marker than other expressions for the purpose of screening and identifying ASD. Future studies can consider the likelihood of using the scanning pattern for angry faces, combined with machine-learning algorithms, to support an ASD diagnosis and improve the accuracy of detecting infants at risk of ASD. Additionally, we need to consider the specificity of markers for ASD. For example, we should tease apart the distinct looking patterns of children with ASD and children with an inhibited or shy temperament in future investigations. An interesting alternative explanation that has been proposed is that eye avoidance of angry faces in children with ASD could be associated with the indirect impact of their greater disorganization of scan patterns caused by emotional arousal. However, this possibility is not testable in our current paradigm. We believe this interesting speculation is a valuable topic for further research and theoretical development.

In conclusion, we observed that 2- to 5-year-old children with ASD tend to look less at others' eyes, especially the eyes of threat-related expressions. We speculate that the diminished eye-looking time for socially threatening expressions may help relieve the threat-elicited hyperarousal caused by direct eye contact. This eye avoidance in ASD could not be explained by reduced face-looking time or enhanced attention or interest to the mouth. Instead, our findings suggest a combination of vigilance to, and avoidance of the core features of threatening faces. Our study also has important implications for the clinical interventions for ASD. Particularly, it suggests that interventions that address social impairments in young children with ASD should incorporate methods of alleviating discomfort elicited by the threatening information of faces. Moreover, the scanning patterns for socially threatening facial expressions can also serve as a potential early marker in developing a computer-aided system to support the diagnosis and early detection of ASD in future investigations.

## References

- Child & Adolescent Psychiatry*, 32, 617–626. <http://dx.doi.org/10.1097/00004583-199305000-00019>
- American Psychiatric Association (APA). (2013). *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, DC: APA Press.
- Bal, E., Harden, E., Lamb, D., Van Hecke, A. V., Denver, J. W., & Porges, S. W. (2010). Emotion recognition in children with autism spectrum disorders: Relations to eye gaze and autonomic state. *Journal of Autism and Developmental Disorders*, 40, 358–370. <http://dx.doi.org/10.1007/s10803-009-0884-3>
- Boll, S., Bartholomaeus, M., Peter, U., Lupke, U., & Gamer, M. (2016). Attentional mechanisms of social perception are biased in social phobia. *Journal of Anxiety Disorders*, 40, 83–93. <http://dx.doi.org/10.1016/j.janxdis.2016.04.004>
- Carré, J. M., McCormick, C. M., & Mondloch, C. J. (2009). Facial structure is a reliable cue of aggressive behavior. *Psychological Science*, 20, 1194–1198. <http://dx.doi.org/10.1111/j.1467-9280.2009.02423.x>
- Chen, Y. P., Ehlers, A., Clark, D. M., & Mansell, W. (2002). Patients with generalized social phobia direct their attention away from faces. *Behaviour Research and Therapy*, 40, 677–687. [http://dx.doi.org/10.1016/S0005-7967\(01\)00086-9](http://dx.doi.org/10.1016/S0005-7967(01)00086-9)
- Corden, B., Chilvers, R., & Skuse, D. (2008). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, 46, 137–147. <http://dx.doi.org/10.1016/j.neuropsychologia.2007.08.005>
- Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., . . . Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nature Neuroscience*, 8, 519–526. <http://dx.doi.org/10.1038/nn1421>
- Dankner, Y., Shalev, L., Carrasco, M., & Yuval-Greenberg, S. (2017). Prestimulus inhibition of saccades in adults with and without attention-deficit/hyperactivity disorder as an index of temporal expectations. *Psychological Science*, 28, 835–850. <http://dx.doi.org/10.1177/0956797617694863>
- De Wit, T. C. J., Falck-Ytter, T., & Hofsten, C. V. (2008). Young children with autism spectrum disorder look differently at positive versus negative emotional faces. *Research in Autism Spectrum Disorders*, 2, 651–659. <http://dx.doi.org/10.1016/j.rasd.2008.01.004>
- Eack, S. M., Mazefsky, C. A., & Minshew, N. J. (2015). Misinterpretation of facial expressions of emotion in verbal adults with autism spectrum disorder. *Autism*, 19, 308–315. <http://dx.doi.org/10.1177/1362361314520755>
- Eisenbarth, H., & Alpers, G. W. (2011). Happy mouth and sad eyes: Scanning emotional facial expressions. *Emotion*, 11, 860–865. <http://dx.doi.org/10.1037/a0022758>
- Falck-Ytter, T., Fernell, E., Gillberg, C., & von Hofsten, C. (2010). Face scanning distinguishes social from communication impairments in autism. *Developmental Science*, 13, 864–875. <http://dx.doi.org/10.1111/j.1467-7687.2009.00942.x>
- Falck-Ytter, T., & von Hofsten, C. (2011). How special is social looking in ASD: A review. *Progress in Brain Research*, 189, 209–222. <http://dx.doi.org/10.1016/B978-0-444-53884-0.00026-9>
- Farabee, D. J., Holcom, M. L., Ramsey, S. L., & Cole, S. G. (1993). Social anxiety and speaker gaze in a persuasive atmosphere. *Journal of Research in Personality*, 27, 365–376. <http://dx.doi.org/10.1006/jrpe.1993.1025>
- Fedor, J., Lynn, A., Foran, W., DiCicco-Bloom, J., Luna, B., & O'Hearn, K. (2017). Patterns of fixation during face recognition: Differences in autism across age. *Autism*. Advance online publication. <http://dx.doi.org/10.1177/13623613171714989>
- Frazier, T. W., Strauss, M., Klingemier, E. W., Zetzer, E. E., Hardan, A. Y., Eng, C., & Youngstrom, E. A. (2017). A meta-analysis of gaze differences to social and nonsocial information between individuals with and without autism. *Journal of the American Academy of Child & Adolescent Psychiatry*, 56, 546–555. <http://dx.doi.org/10.1016/j.jaac.2017.05.005>
- Freeth, M., Chapman, P., Ropar, D., & Mitchell, P. (2010). Do gaze cues in complex scenes capture and direct the attention of high functioning adolescents with ASD? Evidence from eye-tracking. *Journal of Autism and Developmental Disorders*, 40, 534–547. <http://dx.doi.org/10.1007/s10803-009-0893-2>
- Gong, X., Huang, Y., Wang, Y., & Luo, Y. (2011). Revision of the Chinese facial affective picture system. *Chinese Mental Health Journal*, 1, 40–46.
- Gosselin, F., & Schyns, P. G. (2001). Bubbles: A technique to reveal the use of information in recognition tasks. *Vision Research*, 41, 2261–2271. [http://dx.doi.org/10.1016/S0042-6989\(01\)00097-9](http://dx.doi.org/10.1016/S0042-6989(01)00097-9)
- Green, M. J., Williams, L. M., & Davidson, D. (2003). In the face of danger: Specific viewing strategies for facial expressions of threat? *Cognition and Emotion*, 17, 779–786. <http://dx.doi.org/10.1080/026999303002282>
- Hall, J. K., Hutton, S. B., & Morgan, M. J. (2012). Sex differences in scanning faces: Does attention to the eyes explain female superiority in facial expression recognition? *Cognition and Emotion*, 24, 629–637. <http://dx.doi.org/10.1080/02699930902906882>
- Holas, P., Krejtz, I., Cyprianska, M., & Nezelek, J. B. (2014). Orienting and maintenance of attention to threatening facial expressions in anxiety—An eye movement study. *Psychiatry Research*, 220, 362–369. <http://dx.doi.org/10.1016/j.psychres.2014.06.005>
- Horley, K., Williams, L. M., Gonsalvez, C., & Gordon, E. (2003). Social phobics do not see eye to eye: A visual scanpath study of emotional expression processing. *Journal of Anxiety Disorders*, 17, 33–44. [http://dx.doi.org/10.1016/S0887-6185\(02\)00180-9](http://dx.doi.org/10.1016/S0887-6185(02)00180-9)
- Hutt, C., & Ounsted, C. (1966). The biological significance of gaze aversion with particular reference to the syndrome of infantile autism. *Systems Research and Behavioral Science*, 11, 346–356. <http://dx.doi.org/10.1002/bs.3830110504>
- Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder. *Archives of General Psychiatry*, 65, 946–954. <http://dx.doi.org/10.1001/archpsyc.65.8.946>
- Jones, W., & Klin, A. (2013). Attention to eyes is present but in decline in 2–6-month-old infants later diagnosed with autism. *Nature*, 504, 427–431. <http://dx.doi.org/10.1038/nature12715>
- Juth, P., Lundqvist, D., Karlsson, A., & Ohman, A. (2005). Looking for foes and friends: Perceptual and emotional factors when finding a face in the crowd. *Emotion*, 5, 379–395. <http://dx.doi.org/10.1037/1528-3542.5.4.379>
- Kaartinen, M., Puura, K., Mäkelä, T., Rannisto, M., Lemponen, R., Helminen, M., . . . Hietanen, J. K. (2012). Autonomic arousal to direct gaze correlates with social impairments among children with ASD. *Journal of Autism and Developmental Disorders*, 42, 1917–1927. <http://dx.doi.org/10.1007/s10803-011-1435-2>
- Kennedy, D. P., D'Onofrio, B. M., Quinn, P. D., Bölte, S., Lichtenstein, P., & Falck-Ytter, T. (2017). Genetic influence on eye movements to complex scenes at short timescales. *Current Biology*, 27, 3554–3560. <http://dx.doi.org/10.1016/j.cub.2017.10.007>
- Kleinhan, N. M., Richards, T., Weaver, K., Johnson, L. C., Greenon, J., Dawson, G., & Aylward, E. (2010). Association between amygdala response to emotional faces and social anxiety in autism spectrum disorders. *Neuropsychologia*, 48, 3665–3670. <http://dx.doi.org/10.1016/j.neuropsychologia.2010.07.022>
- Kliemann, D., Dziobek, I., Hatri, A., Steimke, R., & Heekeren, H. R. (2010). Atypical reflexive gaze patterns on emotional faces in autism spectrum disorders. *The Journal of Neuroscience*, 30, 12281–12287. <http://dx.doi.org/10.1523/JNEUROSCI.0688-10.2010>
- Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D. (2002). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59, 809–816. <http://dx.doi.org/10.1001/archpsyc.59.9.809>

- Kylliäinen, A., & Hietanen, J. K. (2006). Skin conductance responses to another person's gaze in children with autism. *Journal of Autism and Developmental Disorders*, 36, 517–525. <http://dx.doi.org/10.1007/s10803-006-0091-4>
- Lee, E., Kang, J. I., Park, I. H., Kim, J. J., & An, S. K. (2008). Is a neutral face really evaluated as being emotionally neutral? *Psychiatry Research*, 157, 77–85. <http://dx.doi.org/10.1016/j.psychres.2007.02.005>
- Liu, W., Li, M., & Yi, L. (2016). Identifying children with autism spectrum disorder based on their face processing abnormality: A machine learning framework. *Autism Research*, 9, 888–898. <http://dx.doi.org/10.1002/aur.1615>
- Lord, C., Risi, S., Lambrecht, L., Cook, E. H., Jr., Leventhal, B. L., DiLavore, P. C., . . . Rutter, M. (2000). The autism diagnostic observation schedule-generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30, 205–223. <http://dx.doi.org/10.1023/A:1005592401947>
- Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, 164, 177–190. <http://dx.doi.org/10.1016/j.jneumeth.2007.03.024>
- Matsuda, S., Minagawa, Y., & Yamamoto, J. (2015). Gaze behavior of children with ASD toward pictures of facial expressions. *Autism Research and Treatment*, 2015, 1–8. <http://dx.doi.org/10.1155/2015/617190>
- Mogg, K., & Bradley, B. P. (1998). A cognitive-motivational analysis of anxiety. *Behaviour Research and Therapy*, 36, 809–848. [http://dx.doi.org/10.1016/S0005-7967\(98\)00063-1](http://dx.doi.org/10.1016/S0005-7967(98)00063-1)
- Mogg, K., Garner, M., & Bradley, B. P. (2007). Anxiety and orienting of gaze to angry and fearful faces. *Biological Psychology*, 76, 163–169. <http://dx.doi.org/10.1016/j.biopsycho.2007.07.005>
- Moriuchi, J. M., Klin, A., & Jones, W. (2017). Mechanisms of diminished attention to eyes in autism. *The American Journal of Psychiatry*, 174, 26–35. <http://dx.doi.org/10.1176/appi.ajp.2016.15091222>
- Nakano, T., Tanaka, K., Endo, Y., Yamane, Y., Yamamoto, T., & Nakano, Y. (2010). Atypical gaze patterns in children and adults with autism spectrum disorders dissociated from developmental changes in gaze behaviour. *Proceedings of the Royal Society Biological Sciences*, 277, 2935–2943.
- Nyström, P., Bölte, S., Falck-Ytter, T., & the EASE Team. (2017). Responding to other people's direct gaze: Alterations in gaze behavior in infants at risk for autism occur on very short timescales. *Journal of Autism and Developmental Disorders*, 47, 3498–3509. <http://dx.doi.org/10.1007/s10803-017-3253-7>
- Pan, W., Tian, Y., Wang, X., & Zhang, H. (2018). Ball divergence: Nonparametric two sample test. *Annals of Statistics*, 46, 1109–1137. <http://dx.doi.org/10.1214/17-AOS1579>
- Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual scanning of faces in autism. *Journal of Autism and Developmental Disorders*, 32, 249–261. <http://dx.doi.org/10.1023/A:1016374617369>
- Peltola, M. J., Leppänen, J. M., Vogel-Farley, V. K., Hietanen, J. K., & Nelson, C. A. (2009). Fearful faces but not fearful eyes alone delay attention disengagement in 7-month-old infants. *Emotion*, 9, 560–565. <http://dx.doi.org/10.1037/a0015806>
- Pérez-Edgar, K., Morales, S., LoBue, V., Taber-Thomas, B. C., Allen, E. K., Brown, K. M., & Buss, K. A. (2017). The impact of negative affect on attention patterns to threat across the first 2 years of life. *Developmental Psychology*, 53, 2219–2232. <http://dx.doi.org/10.1037/dev0000408>
- Pierce, K., Müller, R. A., Ambrose, J., Allen, G., & Courchesne, E. (2001). Face processing occurs outside the fusiform 'face area' in autism: Evidence from functional MRI. *Brain: A Journal of Neurology*, 124, 2059–2073. <http://dx.doi.org/10.1093/brain/124.10.2059>
- Quinn, P. C., Yahr, J., Kuhn, A., Slater, A. M., & Pascalis, O. (2002). Representation of the gender of human faces by infants: A preference for female. *Perception*, 31, 1109–1121. <http://dx.doi.org/10.1068/p3331>
- Renzi, C., Tagliaferri, M., & Boehringer, J. (2014). Biases in spatial bisection induced by viewing male and female faces. *Experimental Psychology*, 61, 1–10.
- Rump, K. M., Giovannelli, J. L., Minshew, N. J., & Strauss, M. S. (2009). The development of emotion recognition in individuals with autism. *Child Development*, 80, 1434–1447. <http://dx.doi.org/10.1111/j.1467-8624.2009.01343.x>
- Schurigin, M. W., Nelson, J., Iida, S., Ohira, H., Chiao, J. Y., & Franconeri, S. L. (2014). Eye movements during emotion recognition in faces. *Journal of Vision*, 14, 1–16. <http://dx.doi.org/10.1167/14.13.14>
- Smith, M. L., Cottrell, G. W., Gosselin, F., & Schyns, P. G. (2005). Transmitting and decoding facial expressions. *Psychological Science*, 16, 184–189. <http://dx.doi.org/10.1111/j.0956-7976.2005.00801.x>
- Song, Y., Hakoda, Y., & Sang, B. (2016). A selective impairment in extracting fearful information from another's eyes in Autism. *Autism Research*, 9, 1002–1011. <http://dx.doi.org/10.1002/aur.1583>
- Song, Y., Kawabe, T., Hakoda, Y., & Du, X. (2012). Do the eyes have it? Extraction of identity and positive expression from another's eyes in autism, probed using "Bubbles." *Brain & Development*, 34, 584–590. <http://dx.doi.org/10.1016/j.braindev.2011.09.009>
- Spezio, M. L., Adolphs, R., Hurley, R. S. E., & Piven, J. (2007). Abnormal use of facial information in high-functioning autism. *Journal of Autism and Developmental Disorders*, 37, 929–939. <http://dx.doi.org/10.1007/s10803-006-0232-9>
- Tanaka, J. W., & Sung, A. (2016). The "eye avoidance" hypothesis of autism face processing. *Journal of Autism and Developmental Disorders*, 46, 1538–1552. <http://dx.doi.org/10.1007/s10803-013-1976-7>
- Uljarevic, M., & Hamilton, A. (2013). Recognition of emotions in autism: A formal meta-analysis. *Journal of Autism and Developmental Disorders*, 43, 1517–1526. <http://dx.doi.org/10.1007/s10803-012-1695-5>
- Wang, Q., Hu, C., Short, L. A., & Fu, G. (2012). The influence of shyness on the scanning of own- and other-race faces in adults. *PLoS ONE*, 7, e52203. <http://dx.doi.org/10.1371/journal.pone.0052203>
- Wang, Y., & Luo, Y. (2005). Standardization and assessment of college students' facial expression of emotion. *Chinese Journal of Clinical Psychology*, 13, 396–398.
- Weigelt, S., Koldewyn, K., & Kanwisher, N. (2012). Face identity recognition in autism spectrum disorders: A review of behavioral studies. *Neuroscience and Biobehavioral Reviews*, 36, 1060–1084. <http://dx.doi.org/10.1016/j.neubiorev.2011.12.008>
- Wieser, M. J., Pauli, P., Alpers, G. W., & Mühlberger, A. (2009). Is eye to eye contact really threatening and avoided in social anxiety? An eye-tracking and psychophysiology study. *Journal of Anxiety Disorders*, 23, 93–103. <http://dx.doi.org/10.1016/j.janxdis.2008.04.004>
- Wieser, M. J., Pauli, P., Weyers, P., Alpers, G. W., & Mühlberger, A. (2009). Fear of negative evaluation and the hypervigilance-avoidance hypothesis: An eye-tracking study. *Journal of Neural Transmission*, 116, 717–723. <http://dx.doi.org/10.1007/s00702-008-0101-0>
- Yi, L., Fan, Y., Quinn, P. C., Feng, C., Huang, D., Li, J., . . . Lee, K. (2013). Abnormality in face scanning by children with autism spectrum disorder is limited to the eye region: Evidence from multi-method analyses of eye tracking data. *Journal of Vision*, 13, 227–232. <http://dx.doi.org/10.1167/13.10.5>
- Zhang, Q., Pan, W., Chen, X., and Wang, X. (2018). Ball divergence and its applications to change-point problems for Banach-valued sequences. *Tech. Rep. No. Manuscript in preparation.*

Received June 17, 2017

Revision received May 31, 2018

Accepted June 5, 2018 ■