# Inhibition of Return in the Visual Field

## The Eccentricity Effect is Independent of Cortical Magnification

### Yan Bao,<sup>1,2,\*</sup> Quan Lei,<sup>1,3,\*</sup> Yuan Fang,<sup>1</sup> Yu Tong,<sup>1</sup> Kerstin Schill,<sup>2,4</sup> Ernst Pöppel,<sup>1,2</sup> and Hans Strasburger<sup>2</sup>

<sup>1</sup>Peking University, Beijing, PR China, <sup>2</sup>University of Munich, Germany, <sup>3</sup>Northeastern University, Boston, MA, USA, <sup>4</sup>University of Bremen, Germany

**Abstract.** Inhibition of return (IOR) as an indicator of attentional control is characterized by an eccentricity effect, that is, the more peripheral visual field shows a stronger IOR magnitude relative to the perifoveal visual field. However, it could be argued that this eccentricity effect may not be an attention effect, but due to cortical magnification. To test this possibility, we examined this eccentricity effect in two conditions: the same-size condition in which identical stimuli were used at different eccentricities, and the size-scaling condition in which stimuli were scaled according to the cortical magnification factor (M-scaling), thus stimuli being larger at the more peripheral locations. The results showed that the magnitude of IOR was significantly stronger in the peripheral relative to the perifoveal visual field, and this eccentricity effect was independent of the manipulation of stimulus size (same-size or size-scaling). These results suggest a robust eccentricity effect of IOR which cannot be eliminated by M-scaling. Underlying neural mechanisms of the eccentricity effect of IOR are discussed with respect to both cortical and subcortical structures mediating attentional control in the perifoveal and peripheral visual field.

Keywords: visual attention, inhibition of return, eccentricity effect, cortical magnification, visual field

Effects of visual orienting can be demonstrated by various cueing paradigms. When attention is directed toward a peripheral location by an endogenous cue such as an arrow or a word that predicts where the target would most likely appear, facilitation is typically observed, that is, faster RT for detecting a target at the cued location relative to the uncued location. However, when using an exogenous cue such as a sudden flash of an outline box at a peripheral location, biphasic effects of attentional capture are frequently reported: When the cue-target interval is very brief, observers respond faster to a target at the cued location as compared to an uncued location. When the interval becomes longer, facilitation is reversed to inhibition, that is, observers respond slower to a subsequent target at the cued location. This latter inhibitory effect is typically known as "inhibition of return" (IOR) and has been independently discovered in the mid-1980s by Posner and Cohen (1984) and Tassinari, Aglioti, Chelazzi, Marzi, and Berlucchi (1987). It has been generally assumed that IOR has ecological significance for improving visual search efficiencies: By giving priority to novel locations rather than to those previously attended, visual search efficiency can be greatly improved.

Since the initial discovery of IOR, an increasing amount of research has been dedicated to this phenomenon (Berlucchi, 2006; Klein, 2000; Lupianez, Klein, & Bartolomeo, 2006). Typically, two types of cue-target procedures are employed for its investigation (Bao, Zhou, & Fu, 2004). One is a single-cue procedure, in which a peripheral cue is followed by a target stimulus appearing at either the previously cued location or an opposite, uncued location. The other is a double-cue procedure in which a second cue is added following the peripheral cue so as to reorient attention to the central fixation point before the onset of a following target. Independent of these procedures, IOR has been found to be a very robust phenomenon that can be observed in various situations. It occurs not only in detection tasks but also in discrimination tasks (e.g., Lupianez, Milan, Tornay, Madrid, & Tuleda, 1997; Pratt & Castel, 2001), not only with manual responses but also with eye movement responses (Abrams & Dobkin, 1994), not only in static stimulus displays but also in a dynamic

The first two authors contributed equally to this study.

environment (Tipper, Weaver, Jerreat, & Burak, 1994), and not only in normal young adults but also in other age groups and in patients (Bao et al., 2004; Langley et al., 2007; Larrison-Faucher, Briand, & Sereno, 2002; MacPherson, Klein, & Moore, 2003).

Given the role that IOR might play in the visual search process, it is important to know how IOR is spatially distributed in the visual field. Studies with a method of cueing one peripheral location and measuring the response time (RT) to targets at multiple locations in the visual field suggest that IOR not only dwells at the originally cued location, but also spreads to nearby locations (Maylor & Hockey, 1985) or even to the entire hemi-field with the vertical meridian as a boundary (Berlucchi, Tassinari, Marzi, & Stefano, 1989; Collie, Maruff, Yucel, Danckert, & Currie, 2000; Tassinari et al., 1987). In testing an attentionalmomentum hypothesis with a paradigm of using multiple stimulus locations around a fixation, Pratt and colleagues obtained a different result (Pratt, Adam, & McAuliffe, 1998; Pratt, Spalek, & Bradshaw, 1999). Although the



*Figure 1.* Sample trial sequence of a double-cue IOR paradigm. A valid trial at  $21^{\circ}$  eccentricity is demonstrated for both the same-size condition (a) and the size-scaling condition (b).

demonstrated the eclentricity effect of IOR (Bao & Pöppel, 2007; Bao et al., 2011, 2012, 2013), their lortilal representation was smaller when presented in the periphery than in the perifovea. Therefore, the eccentrility effect of IOR might just reflect lortilal magnifilation, and might not be a valid indicator of a functional dissociation of attentional lontrol in the visual field.

The present study aimed to llarify this possibility by lomparing IOR effects at two stimulus eccentrilities (7° and 21°) in a spatial cueing paradigm lomparing two lritical conditions: one being a same-size condition in which identical stimuli were used for both eccentricities, the other being a size-scaling londition in which the stimuli were scaled according to an estimate of the magnifilation factor. Stimuli were thus larger at 21° than at 7° eccentrility in the latter condition. If cortical magnification is responsible for the eccentricity effect of IOR, then the manipulation of stimulus size (same-size vs. size-scaling londition) should influenle whether the eccentricity effelt of IOR is observed. Specifilally, a larger IOR magnitude at 21° relative to 7° eccentrility would only be expected in the samesize condition and not in the size-slaling condition sinle in the latter the cortical representations of both 7° and 21° targets would be approximately the same.

#### Methods

#### **Participants**

Sixteen university students (10 male and 6 female with an age range of 19–24 years) partilipated in the experiment.

All participants were right-handed and had normal or correlted-to-normal vision. They were all naïve to the purpose of the experiment and were paid after the experiment.

#### Materials and Procedure

The experiment was conducted in a dimly illuminated room. Subjects were seated at 36.5 cm distance from a 19 in. CRT monitor with their heads resting on a lhin rest. The spatial resolution of the CRT monitor was  $1024 \times 768$  pixels and the refresh rate was set at 75 Hz. All stimuli were generated by a PC running Matlab software. Each trial started with a white fixation cross  $(1^{\circ} \times 1^{\circ})$  lentered on the dark lomputer screen. After 1,000 ms, a white outline box serving as a peripheral cue appeared for 100 ms at one of the four peripheral lolations (indicated by dash-line boxes in Figure 1) with equal probability. Two of the peripheral lolations had an eccentrility of 7° (left, right) and the other two an eccentricity of 21° (left, right) (Figure 1). At 70 ms after the offset of the peripheral cue an outline box appeared at the fixation location for 100 ms, reorienting the subject's attention back to the center. Another 70 ms later, a white disk serving as the target appeared at either the same or the symmetric opposite lolation of the peripheral cue, with equal probability. The target remained present until the participant responded, that is, pressed the space bar with the right index finger. Responding was instructed to be as quickly and as accurately as possible. During each trial, subjects were instructed to maintain their fixation at the central cross without shifting their gaze to the peripheral locations. This

Stimulus size	7° Eccentricity			21° Eccentricity		
	Valid	Invalid	IOR	Valid	Invalid	IOR
Same-size	305 (9.4)	282 (10.4)	23 (4.0)	324 (9.9)	287 (10.7)	37 (3.8)
Size-scaling	316 (8.5)	295 (9.3)	21 (5.6)	320 (8.7)	290 (10.4)	30 (5.9)

Table 1. Mean RTs (with Standard Error) for each experimental condition (ms)

requirement was particularly checked by the experimenter during the practice session, and all subjects could follow this instruction successfully before the main test started. An intertrial interval of 1,000 ms with a blank screen was inserted after the subject responded.

The sizes of the cues and targets for different stimulus eccentricities were manipulated as follows: In the same-size condition the cue subtended  $1^{\circ} \times 1^{\circ}$  and the target  $0.6^{\circ} \times 0.6^{\circ}$  for both stimulus eccentricities (7° and 21°); In the size-scaling condition, the peripheral cue at 7° was kept at  $1^{\circ} \times 1^{\circ}$  size while, at 21°, it was magnified to  $2.8^{\circ} \times 2.8^{\circ}$ ; the peripheral target at 7° was kept at  $0.6^{\circ} \times 0.6^{\circ}$  size while, at 21°, it was magnified to  $1.68^{\circ}$ , using a formula for calculating cortical magnification as suggested by Rovamo and Virsu (1979).

The experimental session consisted of 560 trials which were divided into 8 blocks. Four blocks were trials of the same-size condition and 4 blocks were trials of the sizescaling condition. There were 80 catch trials in which no targets were presented and the participants had to withhold their responses. Ten catch trials were randomly mixed with 60 target trials in each block. The number of trials for targets appearing in the left or right visual field, at 7° or 21° eccentricity, and at a cued or uncued location, was balanced in both the same-size and the size-scaling conditions. One half of the participants performed four blocks of the same-size condition first, and the other half performed four blocks of the size-scaling condition first. A practice block with 20 trials of the same-size condition and 20 trials of the size-scaling condition were tested before the main experiment. A short break of approximately 2 min between the experimental blocks was inserted. The entire experimental session lasted approximately 45 min.

#### **Data Preparation**

The total error rate for each subject was lower than 1% and therefore was not further analyzed. Response times for the correct trials of each subject were first submitted to a descriptive analysis. Those RTs outside of 3 standard deviations ( $\pm 3$  *SD*) from the mean were considered as outliers and were excluded from the RT data. A lower limit of 120 ms and a higher limit of 800 ms were further applied to double check whether there remained RTs that were physiologically impossible (too short RTs – less than 120 ms) or were due to reduced concentration (too long RTs – longer than 800 ms). Mean RTs for all experimental conditions are shown in Table 1.

#### Results

Mean RTs were analyzed with a  $2 \times 2 \times 2$  repeated-measure ANOVA, with *Stimulus Size* (same-size and size-scaling), *Stimulus Eccentricity* (7° and 21°), and cue-target *Validity* (valid and invalid) as within-subject variables.

The results of the three-factorial ANOVA showed significant main effects of Stimulus Eccentricity, F(1, 15) =11.308, p < .01,  $\eta_p^2 = .430$ , and cue-target *Validity*, F(1, 15) = 48.153, p < .001,  $\eta_p^2 = .762$ , but no main effect of Stimulus Size, F(1, 15) = 1.492, p > .05,  $\eta_p^2 = .090$ . However, Stimulus Size significantly interacted with Stimulus Eccentricity, F(1, 15) = 31.961, p < .001,  $\eta_p^2 = .681$ . Further analysis of this interaction using paired t-tests with Bonferroni correction showed a slower RT for 21° versus 7° stimuli, but only in the same-size condition (305 ms vs. 294 ms, p < .05), not in the size-scaling condition (305 ms vs. 306 ms, p = .875) (Figure 2a), as expected from the effect of cortical magnification. Consistent with previous observations (Bao & Pöppel, 2007; Bao et al., 2011, 2012, 2013), the interaction between Stimulus Eccentricity and cue-target Validity was also significant, F(1, 15) =33.445, p < .001,  $\eta_p^2 = .690$ . A further paired *t*-test on the magnitude of IOR (i.e., the RT difference between valid and invalid trials) for the two stimulus eccentricities revealed a stronger IOR magnitude for 21° versus 7° stimuli (34 ms vs. 22 ms, t(15) = -5.783, p < .001), showing an eccentricity effect of IOR (Figure 2b). More importantly, no three-way interaction between Stimulus Size, Stimulus *Eccentricity*, and cue-target *Validity* was observed, F(1, 15) = 1.069, p = .318,  $\eta_p^2 = .067$ . This indicated that the eccentricity effect of IOR was not affected by the manipulation of stimulus size. In other words, a result pattern of stronger IOR at 21° versus 7° eccentricity stayed the same in both the same-size condition (37 ms vs. 23 ms) and the size-scaling condition (30 ms vs. 21 ms). No other effects were observed.

#### Discussion

The present study examined whether the difference in cortical projection of the IOR stimuli (as inferred from the cortical magnification factor) is responsible for the eccentricity effect of IOR. By measuring the IOR magnitude at two different stimulus eccentricities ( $7^{\circ}$  and  $21^{\circ}$ ) in a same-size and a size-scaling condition, we observed a robust eccentricity effect of IOR – stronger IOR magnitude for  $21^{\circ}$  relative to  $7^{\circ}$  stimuli – independent of the manipulation of stimulus size.



*Figure 2.* Data illustrating the observed two types of interactions. (a) Interaction between *Stimulus Size* and *Stimulus Eccentricity*. The RT for 21° eccentricity is significantly slower than that for 7° eccentricity in the same-size condition, whereas there is no such difference in the size-scaling condition. (b) Interaction between *Stimulus Eccentricity* and cuetarget *Validity*. A larger IOR magnitude (i.e., RT difference between *valid* and *invalid* trials) for 21° versus 7° eccentricity is observed, that is, an eccentricity effect of inhibition of return. Standard errors are indicated by error bars.

Since in the size-scaling condition, both the cue and the target at the more peripheral location  $(21^\circ)$  were M-scaled simultaneously, we were concerned with a potential confounding of the effects of cue size and target size. Therefore, we further conducted a supplementary experiment with 10 subjects using similar experimental design and procedures, but having only the peripheral target magnified while keeping the peripheral cue size unchanged. Again, we observed a significant eccentricity effect of IOR (i.e., a larger IOR effect for  $21^\circ$  (41 ms) versus 7° (28 ms) eccentricity) which was unaffected by the manipulation of target size.

The main result of this study indicates that cortical magnification is not responsible for the eccentricity effect of IOR, since the critical manipulation in the study – Stimulus Size scaling – did not change the eccentricity effect of IOR. RT was influenced by stimulus size as expected, but IOR, i.e., RT difference, was not. Stimulus size exerts its influence on RT likely by spatial summation of luminous flux up to the size of receptive fields at the respective visual field location (Riccò's Law; Strasburger et al., 2011). The latter can be estimated by a psychophysical procedure known as Westheimer's paradigm (Oehler, 1985). Table 2 shows a comparison of the stimulus sizes used in the present study and estimated mean receptive field sizes at the same location. Stimulus size was just above mean receptive field size at 7° eccentricity, but was below it at 21° in the unscaled condition and above it in the scaled condition. So stimulus sizes compare to summation areas as intended and are thus chosen appropriately.

The observation of a significant two-way interaction between *Stimulus Eccentricity* and cue-target *Validity* in the present study, together with other previous evidence (Bao & Pöppel, 2007; Bao et al., 2011, 2012, 2013), sup-

© 2013 Hogrefe Publishing. Distributed under the Hogrefe OpenMind License http://dx.doi.org/10.1027/a000001 port the notion that the distribution of attention in the human visual field is not homogeneous but can be characterized by (at least) two attentional systems (see also Pöppel & Bao, 2012), one extending from the fovea to approximately  $10^{\circ}$  to  $15^{\circ}$  eccentricity along the horizontal meridian and the other beyond this perifoveal region up to the far periphery.

Although previous studies suggest that IOR is associated with a distributed neural network including the right parietal lobe, superior and middle temporal gyrus, frontal eye field, orbitofrontal cortex, and some subcortical structures including superior colliculus and cerebellum (Lepsien & Pollmann, 2002; Mayer, Seidenberg, Dorflinger, & Rao, 2004; Ro, Farne, & Chang, 2003; Sapir, Soroker, Berger, & Henik, 1999; Vivas, Humphreys, & Fuentes, 2003), it is still too early to specify the neural mechanisms underlying the eccentricity of IOR, since all these previous studies exploring the neural correlates of IOR have only examined the IOR effects in a limited range of stimulus eccentricity, typically within the perifoveal visual field. A direct comparison between IOR in the perifoveal and the peripheral

*Table 2.* Comparison between stimulus size in the present study and mean receptive field size at the two eccentricities used. Mean receptive field sizes are estimated from Equation 18 in Strasburger et al., 2011, based on Oehler (1985)

Eccentricity	Target size	Target size	Receptive
	unsealed	scaled	field size
7°	$0.6^{\circ}$	0.6°	0.42°
21°	$0.6^{\circ}$	1.68°	0.90°

regions of the visual field with fMRI technology comes however from a recent study by Lei and colleagues (2012). Consistent with previous studies, a stronger activation of the dorsal fronto-parietal network was evidenced for the perifoveal IOR, whereas the peripheral IOR was significantly associated with a stronger activation of the prefrontal cortex which indicates that different cortical mechanisms underlie the eccentricity effect of IOR. Due to the limited sensitivity of fMRI in their study, Lei and colleagues could not detect any potential subcortical contributions to the eccentricity effect of IOR.

In a structural model of attentional control (Bao & Pöppel, 2007) it has been argued that different anatomical pathways might be responsible for this functional dissociation. It has been well established that the more peripheral regions of the visual field show a stronger projection to the superior colliculus of the midbrain and from there to extrastriate regions, compared to the foveal and perifoveal regions which possess a strong geniculo-striate projection. Thus, the eccentricity effect of IOR might reflect different attentional control mechanisms which, on the one hand, are dominated by cortical and on the other hand by subcortical processes. In more general terms this functional dissociation of attentional control would highlight the importance of subcortical processing in attention as has been pointed out already some decades ago (Singer, Zihl, & Pöppel, 1977; for review see, e.g., LaBerge, 1995; Sommer & Wurtz, 2004).

Although the eccentricity effect of IOR in the visual field cannot be explained by cortical magnification, stimulus size and eccentricity certainly have an influence on visual processing. If reaction times at 7° and 21° eccentricities were compared for the same-size and the size-scaling conditions, a clear difference was observed. In the samesize condition the more peripheral targets (21°) resulted in significantly longer reaction times compared to the perifoveal targets  $(7^{\circ})$  as also observed in a previous study (Zhou, Bao, Sander, Trahms, & Pöppel, 2010). This difference was no longer present in the size-scaling condition indicating that, indeed, the extent of cortical representation being equalized for the two conditions is relevant for the speed of processing. The fact that the reaction times, after adjusting the size of the stimuli, were similar further strengthens the hypothesis for the eccentricity effect of IOR as not being due to speed differences in perifoveal and peripheral processing. There remains, however, an interesting paradox with respect to the interaction between stimulus size and eccentricity: One might have expected that reaction times for targets at 7° and 21° eccentricities under the size-scaling condition would correspond to the reaction time observed for 7° eccentricity for the same-size condition, but this was not the case; they were the same, but on a higher level corresponding to the slower reaction time under the same-size condition for 21° eccentricity. This result might indicate an instationarity within the experimental conditions as often observed in experiments using reaction time as critical indicator (Pöppel, Schill, Steinbüchel, 1990; Steinbach, Dreden, & Pöppel, 1991). Possibly, when a peripheral target of larger size is used, the overall saliency for all stimuli is increased, resulting in a lower level of general activation which then leads to slower response times. However, whatever may be the case this effect did not compromise the key observation of the eccentricity effect of IOR being independent of the magnification factor. This supports the basic notion of a robust phenomenon of different attentional control in the visual field.

#### Acknowledgment

This research was supported by the National Natural Science Foundation of China (No. 30670703; No. 91120004).

#### References

- Abrams, R. A., & Dobkin, R. S. (1994). Inhibition of return: Effects of attentional cuing on eye movement latencies. *Journal of Experimental Psychology: Human Perception* and Performance, 20, 467–477.
- Bao, Y., & Pöppel, E. (2007). Two spatially separated attention systems in the visual field: Evidence from inhibition of return. *Cognitive Processing*, 8, 37–44.
- Bao, Y., Sander, T., Trahms, L., Pöppel, E., Lei, Q., & Zhou, B. (2011). The eccentricity effect of inhibition of return is resistant to practice. *Neuroscience Letters*, 500, 47–51.
- Bao, Y., Wang, Y., & Pöppel, E. (2012). Spatial orienting in the visual field: A unified perceptual space? *Cognitive Processing*, 13 (Suppl. 1), 93–96.
- Bao, Y., Wang, Z., Liang, W., Wang, Y., Pöppel, E., & Li, H. (2013). Inhibition of return at different eccentricities in the visual field share the same temporal window. *Neuroscience Letters*, 534, 7–11.
- Bao, Y., Zhou, J., & Fu, L. (2004). Aging and the time course of inhibition of return in a static environment. Acta Neurobiologiae Experimentalis, 64, 403–414.
- Bennett, P. J., & Pratt, J. (2001). The spatial distribution of inhibition of return. *Psychological Science*, 12, 76–80.
- Berlucchi, G. (2006). Inhibition of return: a phenomenon in search of a mechanism and a better name. *Cognitive Neuropsychology*, 23, 1065–1074.
- Berlucchi, G., Tassinari, G., Marzi, C. A., & Stefano, M. D. (1989). Spatial distribution of the inhibitory effect of peripheral non-informative cues on simple reaction time to non-fixated visual targets. *Neuropsychologia*, 27, 201–221.
- Collie, A., Maruff, P., Yucel, M., Danckert, J., & Currie, J. (2000). Spatiotemporal distribution of facilitation and inhibitory of return arising from the reflexive orienting of covert attention. *Journal of Experimental Psychology: Human Perception and Performance*, 26, 1733–1745.
- Cowey, A., & Rolls, E. T. (1974). Human cortical magnification factor and its relation to visual acuity. *Experimental Brain Research*, *21*, 447–454.
- Daniel, P. M., & Whitteridge, D. (1961). The representation of the visual field on the cerebral cortex in monkeys. *Journal of Physiology*, 159, 203–221.
- Frost, D., & Pöppel, E. (1976). Different programming modes of human saccadic eye movements as a function of stimulus eccentricity: Indications of a functional subdivision of the visual field. *Biological Cybernetics*, 23, 39–48.
- Klein, R. M. (2000). Inhibition of return. *Trends in Cognitive Science*, 4, 138–147.
- LaBerge, D. (1995). Computational and anatomical models of selective attention in object identification. In M. S. Gazzaniga (Ed.), *The cognitive neurosciences* (1st ed., pp. 649–663). Cambridge, MA: MIT Press.

- Langley, L. K., Fuentes, L. J., Vivas, A. B., & Saville, A. L. (2007). Aging and temporal patterns of inhibition of return. *Journal of Gerontology, Series B: Psychological Sciences* and Social Sciences, 62, 71–77.
- Larrison-Faucher, A., Briand, K. A., & Sereno, A. B. (2002). Delayed onset of inhibition of return in schizophrenia. *Progress in Neuropsychopharmacology & Biological Psychiatry*, 26, 505–512.
- Lei, Q., Bao, Y., Wang, B., & Gutyrchik, E. (2012). fMRI correlates of inhibition of return in perifoveal and peripheral visual field. *Cognitive Processing*, 13 (Suppl. 1), 223–227.
- Lepsien, J., & Pollmann, S. (2002). Covert reorienting and inhibition of return: an event-related fMRI study. *Journal of Cognitive Neuroscience*, 14, 127–144.
- Lupianez, J., Milan, E. G., Tornay, F. J., Madrid, E., & Tuleda, P. (1997). Does IOR occur in discrimination tasks? Yes it does, but later. *Perception & Psychophysics*, 59, 1241–1254.
- Lupianez, J., Klein, R. M., & Bartolomeo, P. (2006). Inhibition of return: Twenty years after. *Cognitive Neuropsychology*, 23, 1003–1014.
- MacPherson, A. C., Klein, R. M., & Moore, C. (2003). Inhibition of return in children and adolescents. *Journal of Experimental Child Psychology*, 85, 337–351.
- Mayer, A. R., Seidenberg, M., Dorflinger, J. M., & Rao, S. M. (2004). An event-related fMRI study of exogenous orienting: Supporting evidence for the cortical basis of inhibition of return? *Journal of Cognitive Neuroscience*, 16, 1262–1271.
- Maylor, E. A., & Hockey, R. (1985). Inhibitory component of externally controlled covert orienting in visual space. *Journal of Experimental Psychology: Human Perception* and Performance, 11, 777–787.
- Oehler, R. (1985). Spatial interactions in the rhesus monkey retina: A behavioural study using the Westheimer paradigm. *Experimental Brain Research*, 59, 217–225.
- Popovic, Z., & Sjöstrand, J. (2001). Resolution, separation of retinal ganglion cells, and cortical magnification in humans. *Vision Research*, 41, 1313–1319.
- Pöppel, E., & Bao, Y. (2012). The visual field paradox: A theoretical account on the reafference principle providing a common frame for the homogeneity and inhomogeneity of visual representation. *Cognitive Processing*, 13 (Suppl. 1), 285–287.
- Pöppel, E., Cramon, D. v., & Backmund, H. (1975). Eccentricity specific dissociation of visual functions in patients with lesions of the central visual pathways. *Nature*, 256, 489–490.
- Pöppel, E., & Harvey, L. O. Jr. (1973). Light-difference threshold and subjective brightness in the periphery of the visual field. *Psychologische Forschung*, 34, 145–161.
- Pöppel, E., Held, R., & Frost, D. (1973). Residual visual function after brain wounds involving the central visual pathways in man. *Nature*, 243, 295–296.
- Pöppel, E., Schill, K., & Steinbüchel, N. v. (1990). Multistable states in intrahemispheric learning of a sensorimotor task. *NeuroReport*, 1, 69–72.
- Posner, M. I., & Cohen, Y. (1984). Components of visual orienting. In H. Bouma & D. G. Bouwhuis (Eds.), Attention and performance X: Control of language processes (pp. 531–556). Hillsdale, NJ: Erlbaum.
- Pratt, J., Adam, J. J., & McAuliffe, J. (1998). The spatial relationship between cues and targets mediates inhibition of return. *Canadian Journal of Experimental Psychology*, 52, 213–216.
- Pratt, J., & Castel, A. D. (2001). Responding to feature or location: A re-examination of inhibition of return and facilitation of return. *Vision Research*, 41, 3903–3908.
- Pratt, J., Spalek, T. M., & Bradshaw, F. (1999). The time to detect targets at inhibited and noninhibited locations: Preliminary evidience for attentional momentum. *Journal* of Experimental Psychology: Human Perception and Performance, 25, 730–746.

- Ro, T., Farne, A., & Chang, E. (2003). Inhibition of return and the human frontal eye fields. *Experimental Brain Research*, 150, 290–296.
- Rovamo, J., & Virsu, V. (1979). An estimation and application of the human cortical magnification factor. *Experimental Brain Research*, 37, 495–510.
- Sapir, A., Soroker, N., Berger, A., & Henik, A. (1999). Inhibition of return in spatial attention: Direct evidence for collicular generation. *Nature Neuroscience*, 2, 1053–1054.
- Schiefer, U., Strasburger, H., Becker, S. T., Vonthein, R., Schiller, J., Dietrich, T. J., & Hart, W. (2001). Reaction time in automated kinetic perimetry: Effects of stimulus luminance, eccentricity, and movement direction. *Vision Research*, 41, 2157–2164.
- Singer, W., Zihl, J., & Pöppel, E. (1977). Subcortical control of visual thresholds in humans: Evidence for modality-specific and retinotopically organized mechanisms of selective attention. *Experimental Brain Research*, 29, 173–190.
- Sommer, M. A., & Wurtz, R. H. (2004). The dialogue between cerebral cortex and superior colliculus: Implications for saccadic target selection and corollary discharge. In L. M. Chalupa & J. S. Werner (Eds.), *The visual neurosciences* (Vol. 2, pp. 1466–1484). Cambridge/MA, London: MIT Press.
- Steinbach, T., Dreden, G. v., & Pöppel, E. (1991). Long-term training in a choice reaction time task reveals different learning characteristics for the visual and auditory system. *Naturwissenschaften*, 78, 185–187.
- Strasburger, H., Rentschler, I., & Harvey, L. O. Jr. (1994). Cortical magnification theory fails to predict visual recognition. *European Journal of Neuroscience*, 6, 1583–1588.
- Strasburger, H., Rentschler, I., & Jüttner, M. (2011). Peripheral vision and pattern recognition: A review. *Journal of Vision*, *11*, 1–82.
- Tassinari, G., Aglioti, S., Chelazzi, L., Marzi, C. A., & Berlucchi, G. (1987). Distribution in the visual field of the costs of voluntarily allocated attention and of the inhibitory after-effects of covert orienting. *Neuropsychologia*, 25, 55–71.
- Tipper, S. P., Weaver, B., Jerreat, L. M., & Burak, A. L. (1994). Object-based and environment-based inhibition of return of visual attention. *Journal of Experimental Psychology: Human Perception and Performance*, 20, 478–499.
- Vivas, A. B., Humphreys, G. W., & Fuentes, L. J. (2003). Inhibitory processing following damage to the parietal lobe. *Neuropsychologia*, 41, 1531–1540.
- Wilson, M. E., & Toyne, M. J. (1970). Retino-tectal and cortical-tectal projections in Macaca mulatta. *Brain Research*, 24, 395–406.
- Zhou, B., Bao, Y., Sander, T., Trahms, T., & Pöppel, E. (2010). Dissociation of summation and peak latencies in visual processing: An MEG study on stimulus eccentricity. *Neuroscience Letters*, 483, 101–104.
- Received October 18, 2012
- Revision received March 20, 2013
- Accepted March 22, 2013
- Published online June 24, 2013

Yan Bao

Department of Psychology Peking University 5 Yiheyuan Road Beijing 100871 PR China Tel. +86 10 62751830 Fax +86 10 62761081 E-mail baoyan@pku.edu.cn