

44

*N. scichinai*

5

*life.scichinai*

\*

10

;

City of We

00871;

Toronto N6A 5L Canada

(29)

[2]

,

mechanisms of

12

[2]

*vision*

,

(ordin masking)  
(surround suppre

.

Po

[29]

,

,

[25]

[27,28]

2

(inner-outer s



[5,50]

(eventually  
calculated potentials,  
fractional magnetic resonance

2.1

(long-range horizonta

6~9

-45]; (

occurred

, V

technique)

[4,2]

[71]

2: 1<sup>[59]</sup>,  
[<sup>[36]</sup>], V4  
0.5 , Bo  
V.  
[77,78],

. Liu  
1/4  
[21]

Chakraborty<sup>[20]</sup>

,

Chakraborty, Cavanagh<sup>[4]</sup>

,

6~8 H, ,

,

Petr

,  
,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

. Fischer<sup>[95]</sup>



fM

,  
Ga  
Gab  
,

a'

[116,117]



M. *alia* [123]

Ga

(perc





on: advantages between line segments. Vision Res, 2007, 47: 437–449. Nat Neurosci, 2001, 43: 171–211.

on und  
f crowd  
isplace  
ical sup  
ntour in  
. Cortic  
ng effect in i  
vision does  
ion with lett  
ontribute to  
ual resolutio  
nines wheth  
ect form vision. J Vi  
cale with target size  
owding. J Vision, 201  
J Vision  
ontralateral  
ankers

and size in a  
wding. Vi  
Ceb  
c

V4 in C  
ive f

g per mination. Later cipher; spat ve.

in visual area V4. *Nature*, 1992, 358:

#### nactivation of the parvocellular

ing the locus of fea-



comple by learning eralize in natural sce  
t al. C cle: natural s and Gestalt rules. P

get selected for recognition: evidence for a common ancestor.

perception, reduce spatial crowding with shrinkage

Perceptual crowding is reduced by binocular cues, such as stereoscopic depth information, which provides information about the three-dimensional position of objects in space.

A typical example of a task involving learning between target and distractors

. Visuals are used during simulations.

W. et al.

Digitized by srujanika@gmail.com

[View Details](#) | [Edit](#) | [Delete](#)

For more information about the study, please contact Dr. Michael J. Hwang at (319) 356-4000 or email at [mhwang@uiowa.edu](mailto:mhwang@uiowa.edu).

in  $\mathbb{R}^n$  is called a *convex function* if and only if

it e ained'

the

increasingly important.

Digitized by srujanika@gmail.com

Digitized by srujanika@gmail.com

www.ijerph.com

For more information about the study, please contact Dr. Michael J. Hwang at (310) 794-3000 or via email at [mhwang@ucla.edu](mailto:mhwang@ucla.edu).